



Actor
portrayal.



A TREATMENT FOR PLATINUM-RESISTANT OVARIAN CANCER

ELAHERE is a prescription medicine used to treat adults with folate receptor-alpha (FR α) positive ovarian cancer, fallopian tube cancer, or primary peritoneal cancer who:

- have not responded to or are no longer responding to treatment with platinum-based chemotherapy **and**
- have received 1 to 3 prior types of chemotherapy.

Your healthcare provider will perform a test to make sure that ELAHERE is right for you. ELAHERE was FDA-approved based on a clinical study that measured how many patients had a tumor response and how long that response lasted. Continued approval is dependent on the results of an ongoing study to confirm the benefit of ELAHERE.

SELECT IMPORTANT SAFETY INFORMATION

What is the most important information I should know about ELAHERE?

ELAHERE can cause serious side effects, including:

Eye problems. Eye problems are common with ELAHERE and can also be severe.

Tell your healthcare provider right away if you develop any eye problems during treatment with ELAHERE, including blurred vision, dry eyes, sensitivity to light, eye pain, or new or worsening vision changes. **Continue reading information about eye problems on page 4.**

Please see Important Safety Information including BOXED WARNING throughout, and [Medication Guide](#).



Actor portrayal.

 **ELAHERE™**
mirvetuximab soravtansine-gynx
injection 100 mg

WHAT SHOULD I ASK MY DOCTOR?



- Have I been tested for FR α , the protein that ELAHERE targets? What were the results?
- Is ELAHERE right for me? (page 4)
- How is it different from other treatments I have been on? (page 6)
- What is the goal of treatment with ELAHERE? (pages 8 and 9)
- What are the side effects I might experience with this treatment? (pages 10 and 12)
- Is there anything I can do to manage those side effects myself? (page 13)
- Will I need to see other specialized doctors? (page 12)
- What can I do to get ready for treatment? (page 15)
- How long will my infusion take? (page 16)
- How often will I have an infusion? (page 16)
- Can I go to my regular eye doctor?
- **Question for your eye doctor:** My oncologist put me on a drug that can affect my eyes. Would your office be able to work with my oncologist to help monitor my eye health during my treatment?

Please see Important Safety Information including BOXED WARNING throughout, and [Medication Guide](#).

IS ELAHERE RIGHT FOR ME?



ELAHERE may be right for you if:



You have platinum-resistant ovarian cancer. This means that it came back or progressed after certain types of chemotherapy



Your cancer is positive for a type of biomarker protein called folate receptor alpha, or FRalpha (FR α)

Before you start treatment, your doctor will test your tissue sample to confirm that your cancer is positive for FR α . To do this, your doctor can use tumor tissue stored from an earlier biopsy or surgery or take a new sample

IMPORTANT SAFETY INFORMATION (CONT'D)

Eye problems. • Your healthcare provider will send you to see an eye care professional to check your eyes before you start treatment with ELAHERE, during treatment with ELAHERE, and as needed for any worsening signs and symptoms of eye problems. • Your healthcare provider will prescribe steroid eye drops and lubricating eye drops before you start and during your treatment with ELAHERE. You should use eye drops as directed by your healthcare provider. • Do not wear contact lenses throughout your treatment with ELAHERE unless you are told to use them by your healthcare provider.

Please see Important Safety Information including BOXED WARNING throughout, and [Medication Guide](#).




Actor portrayal.

 **ELAHERE™**
mirvetuximab soravtansine-gynx
injection 100 mg

HOW DOES ELAHERE KILL CANCER CELLS?



ELAHERE is the first and only FR α -targeted treatment for platinum-resistant ovarian cancer



ELAHERE is a type of treatment called an antibody-drug conjugate (ADC). ADCs are different from traditional chemotherapies and are designed to treat cancer in a targeted way

IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible side effects of ELAHERE? ELAHERE can cause serious side effects, including:

- **Lung problems (pneumonitis).** ELAHERE can cause severe or life-threatening inflammation of the lungs that may lead to death.

Please see Important Safety Information including **BOXED WARNING** throughout, and [Medication Guide](#).



Actor portrayal.

 **ELAHERE™**
mirvetuximab soravtansine-gynx
injection 100 mg

WHAT WERE THE RESULTS OF ELAHERE TREATMENT?



In a clinical study of people who were given ELAHERE*:

~27%

had their tumor shrink by at least 30% after treatment.
This is referred to as the **partial response rate**

~5%

of people had no signs of cancer on a follow-up scan.
This is referred to as the **complete response rate**

~32%

Approximately 1 out of 3 people in the study responded to treatment: their tumor shrank by at least 30% or they had no signs of cancer on a follow-up scan.
This is referred to as the **objective response rate**

*Results from the 104 evaluable people included in the clinical study of 106 people.

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider right away if you get new or worsening symptoms, including trouble breathing, shortness of breath, cough, or chest pain.

Please see Important Safety Information including **BOXED WARNING** throughout, and [Medication Guide](#).

WHAT WERE THE RESULTS OF ELAHERE TREATMENT? (CONT'D)



46.2%

of people in the study achieved **stable disease**. This means their tumors did not grow more than 20% or shrink more than 30%



Median length of time people maintained their response to treatment. A median is the middle number in a group of numbers that are listed from lowest to highest

IMPORTANT SAFETY INFORMATION (CONT'D)

- **Peripheral neuropathy.** You may develop nerve problems called peripheral neuropathy during treatment with ELAHERE. Your healthcare provider will monitor you for signs and symptoms of nerve problems.

Please see Important Safety Information including **BOXED WARNING** throughout, and [Medication Guide](#).

WHAT ARE SOME POSSIBLE SIDE EFFECTS OF ELAHERE?



ELAHERE can cause serious side effects that require medical attention, such as pneumonitis (inflammation of the lungs), peripheral neuropathy (nerve problems), and eye problems. The most common side effects of ELAHERE include:

- Feeling tired
- Increased liver enzymes in the blood
- Nausea
- Stomach-area (abdominal) pain
- Decreased red or white blood cell counts
- Diarrhea
- Decreased albumin level in the blood
- Constipation
- Decreased magnesium level in the blood

Due to side effects: 11% of people stopped treatment, 39% of people had dose delays, 20% of people had their dose reduced, and 31% of people reported serious adverse events.

These are not all the possible side effects of ELAHERE. Call your doctor for medical advice about side effects. You may report side effects to the FDA at **1-800-FDA-1088**

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider if you get new or worsening numbness or tingling in your hands or feet or muscle weakness.

Please see Important Safety Information including **BOXED WARNING** throughout, and [Medication Guide](#).



Actor portrayal.



ELAHERE™
mirvetuximab soravtansine-gynx
injection 100 mg

WILL ELAHERE AFFECT MY EYES?



Some people may develop eye problems during treatment with ELAHERE. These may include blurred vision, dry eyes, sensitivity to light, eye pain, or new or worsening vision changes. Tell your doctor if you have a history of vision or eye problems.



- Eye-related side effects most commonly occurred between the second and third infusion of ELAHERE
- **Less than 1% of people in the study stopped treatment because of eye problems**
- In clinical studies, no patients had permanent eye-related side effects

Your doctor will also have you meet with an eye doctor (ophthalmologist or optometrist) before your first infusion and periodically throughout your treatment. You will receive eye drops to use during your treatment

IMPORTANT SAFETY INFORMATION (CONT'D)

The most common side effects of ELAHERE include: • feeling tired • increased liver enzymes in the blood • nausea • stomach-area (abdominal) pain • decreased red or white blood cell counts • diarrhea • decreased albumin level in the blood • constipation • decreased magnesium level in the blood

Your healthcare provider may change your dose of ELAHERE, delay treatment, or completely stop treatment if you have side effects.

Please see Important Safety Information including **BOXED WARNING** throughout, and [Medication Guide](#).


HOW SHOULD I CARE FOR MY EYES?



To help manage your risk, here are steps you can take:



- Your doctor will prescribe steroid eye drops and over-the-counter, lubricating, preservative-free eye drops for use before you start treatment, and during your treatment, with ELAHERE
- You should use eye drops as directed by your doctor. Refer to the ELAHERE ocular management brochure for more detailed information
- Avoid wearing contact lenses throughout your treatment with ELAHERE unless your doctor tells you that you can
- **During treatment with ELAHERE, tell your doctor if you experience any eye problems, including blurred vision, dry eyes, sensitivity to light, eye pain, or new or worsening vision changes**



Need reminders to use your eye drops? There are several mobile apps available for download, designed to help you and/or your caregiver remember your eye drop schedule. You or your caregiver can find a variety to choose from at your preferred app store

IMPORTANT SAFETY INFORMATION (CONT'D)

What should I tell my healthcare provider before receiving ELAHERE? Tell your healthcare provider about all of your medical conditions, including if you:

- have vision or eye problems.
- have liver problems.
- are pregnant or plan to become pregnant.

Please see Important Safety Information including BOXED WARNING throughout, and [Medication Guide](#).



Actor portrayal.

 **ELAHERE™**
mirvetuximab soravtansine-gynx
injection 100 mg

WHAT HAPPENS BEFORE I START ELAHERE?



FRα

Before you start treatment, your doctor will test a sample from your tumor to confirm that your cancer is FRα positive. Your doctor may be able to use tissue from an earlier biopsy or surgery for testing

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking ELAHERE with certain other medicines may cause side effects.

- Before each dose of ELAHERE, you will receive medicines to help prevent infusion-related reactions, nausea, and vomiting
- Some of these medications may be taken orally and some may be infusions, which are delivered through your vein. A needle or catheter will be used to start the infusion
- Usually, these medications will be given on the same day as the infusion. However, your doctor may choose to give you the medications the day before you start ELAHERE

IMPORTANT SAFETY INFORMATION (CONT'D)

ELAHERE can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ELAHERE.

Please see Important Safety Information including BOXED WARNING throughout, and [Medication Guide](#).

HOW WILL I RECEIVE ELAHERE?



ELAHERE is given by infusion (IV or a port) every 3 weeks

**Typical
hours per
infusion**



1.5 to 3

...



1 to 2

...



1 to 2

...

Your doctor will
decide how
many treatments
you need

ELAHERE dosage is determined by your weight. To ensure accurate dosing, your doctor will weigh you prior to treatment.

If you have side effects with ELAHERE, your doctor may reduce your dose, delay your dose, or completely stop treatment

IV=intravenous.

IMPORTANT SAFETY INFORMATION (CONT'D)

Females who are able to become pregnant: • Your healthcare provider should do a pregnancy test before you start treatment with ELAHERE. • You should use an effective birth control (contraception) during treatment and for 7 months after your last dose of ELAHERE.

Please see Important Safety Information including **BOXED WARNING** throughout, and [Medication Guide](#).



Actor portrayal.

 **ELAHERE™**
mirvetuximab soravtansine-gynx
injection 100 mg

IS THERE FINANCIAL SUPPORT FOR ELAHERE?




ELAHERE Support Services: Here to help you get started on ELAHERE

Once enrolled in the program, your ELAHERE Support Services Case Manager will help you understand your insurance coverage and navigate financial assistance options.



For patients with commercial insurance*:

**Eligible patients could pay
as little as \$0 for their medication**



Patient Assistance Program (PAP)

The Patient Assistance Program (PAP) provides eligible patients with ELAHERE at no charge[†]

*Terms and conditions apply. Patients are eligible for co-pay assistance if enrolled in private commercial health insurance and are not covered by state or federal healthcare programs, and who meet the eligibility criteria. Patients will be enrolled for 12 months. There are no income requirements to participate in the program.

[†]Criteria include: patients who are uninsured or have insurance that excludes coverage for ELAHERE (including patients on Medicare or Medicaid), residents of the United States or Puerto Rico, and patients who meet the financial eligibility requirements. Terms and conditions apply.

IMPORTANT SAFETY INFORMATION (CONT'D)

- are breastfeeding or plan to breastfeed. It is not known if ELAHERE passes into your breast milk. Do not breastfeed during treatment and for 1 month after your last dose of ELAHERE.

Please see Important Safety Information including BOXED WARNING throughout, and [Medication Guide](#).

IS THERE FINANCIAL SUPPORT FOR ELAHERE? (CONT'D)



Visit www.ELAHERE.com for more information on the ELAHERE Support Services Program and for frequently asked questions



Questions? Connect with an ELAHERE Support Services Program specialist

Phone: 1-833-ELAHERE (1-833-352-4373) Monday to Friday, 8 AM to 8 PM ET

Email: ELAHERESupport@cardinalhealth.com

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking certain other medicines during treatment with ELAHERE may cause side effects.

Please see Important Safety Information including BOXED WARNING throughout, and [Medication Guide](#).

WHAT ELSE SHOULD I DISCUSS WITH MY DOCTOR?



Before receiving ELAHERE, tell your doctor about all of your medical conditions, including if you:

- Have vision or eye problems.
- Have liver problems.
- Are pregnant or plan to become pregnant. ELAHERE can harm your unborn baby. Tell your doctor right away if you become pregnant or think you may be pregnant during treatment with ELAHERE.
- Are breastfeeding or plan to breastfeed. It is not known if ELAHERE passes into your breast milk. Do not breastfeed during treatment and for 1 month after your last dose of ELAHERE.

If you are able to become pregnant:

- Your doctor should do a pregnancy test before you start treatment with ELAHERE
- You should use an effective birth control (contraception) during treatment and for 7 months after your last dose of ELAHERE

IMPORTANT SAFETY INFORMATION (CONT'D)

These are not all of the possible side effects of ELAHERE. Call your doctor for medical advice about side effects. You are encouraged to report side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see Important Safety Information including BOXED WARNING throughout, and [Medication Guide](#).



Actor portrayal.

 **ELAHERE™**
mirvetuximab soravtansine-gynx
injection 100 mg



Actor
portrayal.



SUPPORT THROUGHOUT YOUR TREATMENT JOURNEY



We're here to help. Contact us to learn about
our support programs and financial assistance.

For questions, call **1-833-ELAHERE**
Monday-Friday, 8:00 AM to 8:00 PM ET
Or email **ELAHERESupport@cardinalhealth.com**

Please see Important Safety Information including **BOXED WARNING** throughout,
and **Medication Guide**.

immun•gen

© 2022 ImmunoGen, Inc., All Rights Reserved. US-MIRV-2200089v4.0 04/2023
ELAHERE and the ELAHERE logo are trademarks or registered trademarks of ImmunoGen, Inc.
ImmunoGen and the ImmunoGen logos are trademarks or registered trademarks of ImmunoGen, Inc.