

Category

Best Digital Health Solution

General Information**Company Name ***

WARD 24/7

Number of employees *

11-50

Turnover and/or Funding

2019 - Public grant \$7,500,000

2020 - Pre-seed = \$600,000

2022/2023 - Seed A 2 = \$3.000.000

2024/2025 - Seed B 3 = \$3.500.000

words remaining :

476

Product/Solution Name *

WARD Clinical Support System (WARD-CSS)

Corporate Name *

WARD247 ApS

Date of Approval *

2025-02-20

Indications *

WARD-CSS is indicated for use for patients, 18 years and older, and intended for but not restricted to, patients at risk for critical complications. WARD-CSS notifies healthcare professionals and indicates that attention should be brought to the patient when deviating vital signs are detected. WARD-CSS is an adjunct to, and is not intended to replace, existing vital sign monitoring practices but alerts the health care professionals to changes in the patient's physiological status.

words remaining :

427

Therapeutic Areas *

WARD-CSS provides physiological insight into a patient's risk of critical complications related to:

- Bradypnea

- Tachypnea
- Desaturation and bradypnea
- Desaturation
- Tachycardia
- Bradycardia
- Hypotension
- Hypertension
- Hypotension and tachycardia
- Hypotension and bradycardia
- Fever

words remaining :

459

*Kindly clearly label your files with company name and asset name.

Background information and need for drug / device

(please be as specific as possible in your description; limit 500 words)

Each year, tens of thousands of hospitalized patients experience preventable complications, many of which are preceded by deviations in vital signs that go unnoticed due to infrequent or inefficient monitoring practices.

Current standard of care on general wards involves periodic spot checks, which often fail to detect early signs of deterioration.

The challenge is further exacerbated by global nurse shortages, alarm fatigue, and the lack of scalable, intelligent monitoring solutions.

WARD 24/7 was founded by clinicians to address this exact need.

The WARD-CSS platform transforms general ward care by enabling continuous vital sign monitoring via wireless sensors, providing clinically relevant alerts, and allowing for timely interventions that improve patient outcomes and optimize clinical resources.

words remaining :

386

*Kindly clearly label your files with company name and asset name.

History of the development of the solution/product *

(please be as specific as possible in your description; 500 words)

Research history

2019

- 2.300 patients observed
- 2 large RCTs (700 patients)
- 6 ongoing studies
- 23 peer reviewed publications
- 6 PhDs

2020

- Prototype
- Licence-agreement
- Patent pending

2022/2023

- MDR (CE) certification
- Product launch (WARD-CSS)
- First sale
- EU & US pilots
(including
Cleveland Clinic)

2024/2025

- Strong US partnerships
- Sensor partners
- EU launch
- Gen 2 prototype

The problem

Hospitalized patients often get complications:

30% unexpected complications

40% of those are potentially preventable

70% of mortality after complications happens in the general wards

4-6 days prolonged stay in hospital after complications

\$10,500 - \$15,000 added costs for a patient with moderate / severe complications

words remaining :

385

*Kindly clearly label your files with company name and asset name.

Attached Files:

- [WARD247 MEDICAL CLAIMS DECK WARDCSS V13x.pptx](#)
- [WARD247 MEDICAL CLAIMS DECK WARDCSS V13x.pptx](#)

Why this drug or device is innovative, the broad implications for future research, and/or how it will improve the human condition *

WARD-CSS is a software as a medical device (SaMD) that integrates wireless continuous monitoring with clinically validated algorithms, providing:

- Detects clinical deterioration 48 hours earlier than standard monitoring ($p < 0.05$)
- Reduces non-actionable alerts by over 80% ($p < 0.05$)
- Operates with >96% sensitivity for sustained deviations
- Interfaces with existing hospital systems and is sensor-agnostic

Unlike generic monitoring or predictive analytics tools, WARD-CSS provides clear, clinically actionable alerts, reducing nurse workload and alert fatigue while maintaining high sensitivity and safety.

words remaining :

420

*Kindly clearly label your files with company name and asset name.

Please provide appropriate references (PubMed, Abstract, Website) *

Logo

Description automatically generated, Picture

The references are listed as what references there is on the different pages/slides in the PPTX

P. 2,3,5,6,7,8,9,10

Source: Clinical Evaluation Report

P. 12

(2) Ghaferi AA, Birkmeyer JD, Dimick JB. Variation in hospital mortality associated with inpatient surgery. N Engl J Med 2009;361:1368-75.

An estimation of the global volume of surgery: a modelling strategy based on available data. July 2008, The Lancet. T G Weiser et al.
Br J Anaesth 2016;117:601-9.

Link: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(08\)60878-8/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(08)60878-8/fulltext)

3rd leading cause of death (controversial claim)
Nepogodiev D, Martin J, Biccard B, m.fl.
Global burden of postoperative death. Lancet 2019. s. 401.

Link:

Devereaux PJ, Sessler DI. Cardiac Complications in Patients Undergoing Major Noncardiac Surgery. N Engl J Med 2015;373:2258-69.

Global patient outcomes after elective surgery: prospective cohort study in 27 low-, middle- and high-income countries. The International Surgical Outcomes Study group. Br J Anaesth. 2016

P. 14

(7) Data visualized from: PROVE Network Investigators for the Clinical Trial Network of the European Society of Anaesthesiology, Hemmes SN, Gama de Abreu M, Pelosi P, Schultz MJ. High versus low positive end-expiratory pressure during general anaesthesia for open abdominal surgery (PROVHILO trial): a multicentre randomised controlled trial. Lancet. 2014;384(9942):495-503. doi:10.1016/S0140-6736(14)60416-5

Link: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(14\)60416-5/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(14)60416-5/fulltext)

P. 15,17,18,24

(8) Elvekjaer M, Rasmussen SM, Grønbæk KK, m.fl. Clinical impact of vital sign abnormalities in patients admitted with acute exacerbation of chronic obstructive pulmonary disease: an observational study using continuous wireless monitoring. Intern Emerg Med [Internet] 2022; Available from: <http://dx.doi.org/10.1007/s11739-022-02988-w>

P 15,17

(9) Continuous monitoring of vital sign abnormalities; association to clinical complications in 500 postoperative patients. Haahr-Raunkjaer C, Mølgaard J, Elvekjaer M, Rasmussen SM, Achiam MP, Jorgensen LN, Søgaard MIV, Grønbaek KK, Oxbøll AB, Sørensen HBD, Meyhoff CS, Aasvang EK. Acta Anaesthesiol Scand. 2022 May;66(5):552-562. doi:10.1111/aas.14048. Epub 2022 Feb 28.

P. 16

(10) Complications in Surgical Patients, JAMA surgery 2002, Mark A. Healey et al

P. 18

(11) Incidence and type of complications in non-operated patients at a surgical ward, Eelco J Veen et al, 2010

(12) Risk factors, costs and complications of delayed hospital discharge from internal medicine wards at a Canadian academic medical centre: retrospective cohort study, Anthony D. Bai

P. 19

(13) Data visualized from

Risk of mortality if admission with 1 or more vital sign deviations.

Prognosis and Risk Factors for Deterioration in Patients Admitted to a Medical Emergency Department. PLoS One 2014, Henriksen DP et al..

P. 20,23

(14) Association between clinically abnormal observations and subsequent in-hospital mortality: a prospective study, Resuscitation 2004, Michael Buist et al

Link: [https://www.resuscitationjournal.com/article/S0300-9572\(04\)00123-6/fulltext](https://www.resuscitationjournal.com/article/S0300-9572(04)00123-6/fulltext)

P. 21

(15) An early warning scoring system for detecting developing critical illness RJM Morgan, F Williams, MM Wright - Clin Intensive Care, 1997

- not available

(16) Royal College of Physicians. National Early Warning Score (NEWS): Standardising the Assessment of Acute Illness Severity in the NHS. Report of a Working Party.

(17) Bryan Williams. The National Early Warning Score: from concept to NHS implementation. DOI: 10.7861/clinmed.2022-news-concept

Link: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9761416/>

(18) Smith GB, Prytherch DR, Schmidt PE, Featherstone PI. Review and performance evaluation of aggregate weighted 'track and trigger' systems. *Resuscitation*. 2008;77:170-9.

Link: [https://www.resuscitationjournal.com/article/S0300-9572\(07\)00653-3/fulltext](https://www.resuscitationjournal.com/article/S0300-9572(07)00653-3/fulltext)

(19) Subbe CP, Slater A, Menon D, Gemmell L. Validation of physiological scoring systems in the accident and emergency department. *Emerg Med J*. 2006;23:841-5.

Link: <https://emj.bmj.com/content/23/11/841>

(20) Groarke JD, Gallagher J, Stack J, Aftab A, Dwyer C, McGovern R, et al. Use of an admission early warning score to predict patient morbidity and mortality and treatment success. *Emerg Med J*. 2008;25:803-6.

Link: <https://emj.bmj.com/content/25/12/803>

(21) Cretikos M, Chen J, Hillman K, Bellomo R, Finfer S, Flabouris A. The objective medical emergency team activation criteria: A case-control study. *Resuscitation*. 2007;73:62-72.

Link: [https://www.resuscitationjournal.com/article/S0300-9572\(06\)00564-8/fulltext](https://www.resuscitationjournal.com/article/S0300-9572(06)00564-8/fulltext)

(22) Barlow GD, Nathwani D, Davey PG. Standardised early warnings scoring system. *Clin Med*. 2006;6:422-3. author reply 423-424.

(23) Paterson R, MacLeod DC, Thetford D, Beattie A, Graham C, Lam S, et al. Prediction of in-hospital mortality and length of stay using an early warning scoring system: Clinical audit. *Clin Med*. 2006;6:281-4.

(24) Subbe CP, Gao H, Harrison DA. Reproducibility of physiological track-and-trigger warning systems for identifying at-risk patients on the ward. *Intensive Care Med*. 2007;33:619-24.

(25) Bhatnagar M, Sirohi N, Dubey AB. Prediction of hospital outcome in emergency medical admissions using modified early warning score (MEWS): Indian experience. *J Family Med Prim Care*. 2021 Jan;10(1):192-198. doi: 10.4103/jfmpc.jfmpc_1426_20. Epub 2021 Jan 30. PMID: 34017725; PMCID: PMC8132807.

(26) Nakitende I, Nabiryo J, Namujwiga T, Wasingya-Kasereka L, Kellett J. Do different patient populations need different early warning scores? The performance of nine different early warning scores used on acutely ill patients admitted to a low-resource hospital in sub-Saharan Africa. *Clin Med (Lond)*. 2020 Jan;20(1):67-73. doi: 10.7861/clinmed.2019-0196. Epub 2019 Nov 8. PMID: 31704729; PMCID: PMC6964175.

P. 21,35

(27) Gao H, McDonnell A, Harrison DA, Moore T, Adam S, Daly K, et al. Systematic review and evaluation of physiological track and trigger warning systems for identifying at-risk patients on the ward. *Intens Care Med*. 2007;33:667-79.

P. 22

(28) Nielsen et al. Review of Early Warning Score in preventing sudden critical illness and death.

Link: <https://pubmed.ncbi.nlm.nih.gov/30327089/>

(29) Gerry et al. Early warning scores for detecting deterioration in adult hospital patients: systematic review and critical appraisal of methodology. *BMJ* 2020;369:m150.

P. 22,23

(30) Early warning systems and rapid response systems for the prevention of patient deterioration on acute adult hospital wards, Jennifer McGaughey, Dean A Fergusson, Peter Van Bogaert, and Louise Rose

P. 23

(31) Kause et al: A comparison of Antecedents to Cardiac Arrests, Deaths and Emergency Intensive care Admissions in Australia and New Zealand, and the United Kingdom - The ACADEMIA study

Link: [https://www.resuscitationjournal.com/article/S0300-9572\(04\)00247-3/fulltext](https://www.resuscitationjournal.com/article/S0300-9572(04)00247-3/fulltext)

P. 25

(32) Hypoxemia Within the First 3 Postoperative Days Is Associated With Increased 1-Year Postoperative Mortality After Adjusting for Perioperative Opioids and Other Confounders, ANESTHESIA & ANALGESIA 2019, Karsten Bartels, et al.

(33) Perioperative myocardial injury and the contribution of hypotension, Intensive Care Medicine, 2018, Daniel I. Sessler & Ashish K. Khanna

P. 26

(34) Grønbæk et al. Continuous monitoring is superior to manual measurements in detecting vital sign deviations in patients with COVID-19. Acta Anaesthesiol Scand. 2023 May;67(5):640-648

P. 27

(35) Farenden, Gamble, Welch: Impact of implementation of the National Early Warning Score on patients and staff. British journal of hospital medicine (London, England: 2005) 78(3):132-136

Link: <https://pubmed.ncbi.nlm.nih.gov/28277766/>

P. 28

(36) Clifton DA, Clifton L, Sandu DM, Smith GB, Tarassenko L, Vollam SA, et al. "Errors" and omissions in paper-based early warning scores: the association with changes in vital signs-a database analysis. BMJ Open. 2015;5:1-7

Link: https://www.researchgate.net/publication/280030418_27Errors%27_and_omissions_in_paper-based_early_warning_scores_The_association_with_changes_in_vital_signs-a_database_analysis

(37) Jansen JO, Cuthbertson BH. Detecting critical illness outside the ICU: the role of track and trigger systems. *Curr Opin Crit Care*. 2010;16:184-90. <https://doi.org/10.1097/MCC.0b013e328338844e>

Link: https://journals.lww.com/co-criticalcare/abstract/2010/06000/detecting_critical_illness_outside_the_icu__the.3.aspx

P. 28,38

(38) Areia C, Biggs C, Santos M, m.fl. The impact of wearable continuous vital sign monitoring on deterioration detection and clinical outcomes in hospitalised patients: a systematic review and meta-analysis. *Crit Care* 2021;25:351.

P. 29,30,31,32

(39) Redfern OC, Griffiths P, Maruotti A, m.fl.

The association between nurse staffing levels and the timeliness of vital signs monitoring: a retrospective observational study in the UK.

BMJ Open British Medical Journal Publishing Group; 2019;9:e032157.

P. 33

(40) Pedersen NE, Rasmussen LS, Petersen JA, m.fl.

A critical assessment of early warning score records in 168,000 patients.

J Clin Monit Comput 2018;32:109-16.

P. 34

(41) <https://nyheder.tv2.dk/2021-10-14-sygeplejestuderende-blev-sat-til-at-passe-en-doeende-patient-dagen-efter-kom-beskeden-hun>

(42) <https://omtv2.tv2.dk/nyheder/2021/10/tv-2-dokumentar-opraab-fra-sygehuset>

P. 36

(43) Mariska Weenk, MD, et al. Continuous Monitoring of Vital Signs in the General Ward Using Wearable Devices: Randomized Controlled Trial, JOURNAL OF MEDICAL INTERNET RESEARCH, 2020

P. 37

(44) Reducing ICU Utilization, Length of Stay, and Cost by Optimizing the Clinical Use of Continuous Monitoring System Technology in the Hospital, Patricia C. Dykes et al.

Link: [https://www.amjmed.com/article/S0002-9343\(21\)00698-7/fulltext](https://www.amjmed.com/article/S0002-9343(21)00698-7/fulltext)

P. 39

(45) Siebig, S. , Kuhls, S. , Imhoff, M. , Gather, U. , Schölmerich, J. & Wrede, C. E. (2010). Critical Care Medicine, 38 (2), 451-456. doi: 10.1097/CCM.0b013e3181cb0888.

Link:

https://journals.lww.com/ccmjournal/abstract/2010/02000/intensive_care_unit_alarms_how_many_do_w

(46) Bonafide CP, Lin R, Zander M, et al. Association between exposure to nonactionable physiologic monitor alarms and response time in a children's hospital. J Hosp Med. 2015;10(6):345-351. doi:10.1002/jhm.2331

(47) Drew BJ, Harris P, Zègre-Hemsey JK, Mammone T, Schindler D, Salas-Boni R, Bai Y, Tinoco A, Ding Q, Hu X. Insights into the problem of alarm fatigue with physiologic monitor devices: a comprehensive observational study of consecutive intensive care unit patients. PLoS One. 2014 Oct 22;9(10):e110274. doi:10.1371/journal.pone.0110274. PMID: 25338067; PMCID: PMC4206416.

(48) Atzema, Clare et al. "ALARMED: adverse events in low-risk patients with chest pain receiving continuous electrocardiographic monitoring in the emergency department. A pilot study." The American journal of emergency medicine vol. 24,1 (2006): 62-7

Link: https://www.researchgate.net/publication/7430302_ALARMED_Adverse_events_in_Low-risk_patients_with_chest_pain_Receiving_continuous_electrocardiographic_Monitoring_in_the_Emerge

P. 40

(49) Monitor Alarm Fatigue: An Integrative Review, Biomedical instrumentation and technology, 2012, Maria Cvach

P. 41

(50) Feasibility of wireless continuous monitoring of vital signs without using alarms on a general surgical ward: A mixed methods study, PLOS one 2022, Jobbe P.L. Leenen et al

P. 44

(51) Unpublished time study, in process to be published, available on demand

P. 44,57,58,59

(52) Mølgaard et al Preventing patient vital sign deterioration in post surgical patients using wireless observation and a modified staff alerting algorithm - A pragmatic, randomized controlled trial. 2023 - in preparation

- not available

P. 44

(53) Based on Cost of SAEs from Cost of adverse events: Kjellberg et al. BMC Health Services Research (2017) 17:651

DOI 10.1186/s12913-017-2605-5 - applying reduction in clinical complications from unpublished RCT Sample Text

Link: <https://bmchealthservres.biomedcentral.com/articles/10.1186/s12913-017-2605-5>

P. 46

(54) C. L. Duuset al.

Continuous vital sign monitoring after major abdominal surgery- Quantification of micro events

Acta Anaesthesiol Scand. 2018.

P. 47,48,49

(55) Own unpublished analysis. Data from

Vang Jensen et al - Early detection of clinical complications by wireless continuous

monitoring with AI-alerts compared to clinical practice - in preparation

. not available

P. 50

(56)

<https://ward247.atlassian.net/wiki/spaces/AI/pages/434798604/Alert+functionality+frequency+and+distril>

P. 51,53,54

(57) Own unpublished analysis. Data from

Kjærsgaard et al Using technical filtering and clinical criteria reduces vital sign alerts using continuous monitoring in the ward - in preparation

. not available

P. 52

(58) Sigvardt et al. Workload associated with manual assessment of vital signs using the National Early Warning Score as compared with continuous wireless monitoring - In preparation

. not available

P. 55

(59) <https://ward247.atlassian.net/wiki/spaces/EQ/pages/442302578/Summative+evaluation+WARD-CSS>

P. 56

(60) SAEs cause 4 days prolonged stay in hospitals

Direct medical costs of adverse events in Dutch hospitals, BMC Health Services Research volume 9, Article number: 27 (2009) Lilian HF Hoonhout et al.

(61) 3 worst clavien-dindorated complications cost 9.505 EUR during primary admission

Costs associated with adverse events among acute patients. BMC Health Services Research 2017. Jakob Kjellberg

*Kindly clearly label your files with company name and asset name.